

Books at Amazon.com on Nanotechnology

<http://www.amazon.com/s/103-9040246-0256655?ie=UTF8&keywords=NANOTECHNOLOGY%20&tag=thenanotecinscie&index=books&Search=Search&link%5Fcode=qs>

Ions

Ions are charged species.

This is due to a gain or loss of electrons (almost never protons!)



Gain of electrons → negatively charged ion ≡ **anion**

Loss of electrons → positively charged ion ≡ **cation**

There are two types of ions: simple and complex

Simple ions

Simple ions (monoatomic ions) are species with one nucleus that has gained or lost electrons



Complex ions

Complex ions are molecules, where the atoms are connected by covalent bonds, that have lost or gained electrons.

Those are also called polyatomic ions

Examples: NH_4^+ ammonium ion or H_3O^+ hydronium ion

Note that polyatomic ions themselves do not break apart

More examples can be found in **Table 2.2 of M & H p. 41**

Compounds can also be formed the interaction of anions and cations (species of opposite charge)

These are called ionic compounds even though they are electrically neutral (# of positive charges = # negative charges)

Such compounds are held together by an electrostatic attraction known as an ionic bond

Crystalline table salt (NaCl) is Na^+ and Cl^- bonded together. When dissolved, NaCl dissociates to form a solution of Na^+ and Cl^-

Names of Compounds

A compound can be identified using its formula (NaCl) or its name (sodium chloride).

The following are conventions used in naming compounds:

1. Ions

Simple (monoatomic) cations of metals take the name of the metal

Na⁺ = sodium K⁺ = potassium

However, some metals, specially transition metals in the periodic table, exist in more than one cationic form.

To distinguish these, we note the charge as a Roman numeral

Fe²⁺ = Fe(II) Fe³⁺ = Fe(III)

Simple (monoatomic) anions are named by adding the suffix **ide** to the base name of the non-metal

Cl⁻ = chloride O²⁻ = oxide H⁻ = hydride

Polyatomic ions are given special names (see **M&H Table 2.2**).

However, those containing oxygen (oxoanions) can be named using the base name and the following rules:

When the non-metal forms only two oxoanions, the suffix **ate** is used for the one with the more oxygens.

The suffix **ite** is used for the one with the least oxygens

SO₄²⁻ = sulfate SO₃²⁻ = sulfite

When a non-metal forms more than two oxoanions, we use the prefixes, **per** and **hypo** for the **most** and **least** oxygens, respectively

ClO_4^- = perchlorate

ClO_3^- = chlorate

ClO_2^- = chlorite

ClO^- = hypochlorite

2. Ionic Compounds

Recall that ionic compounds consist of a cation and an anion. It follows that they are named using the names of the cation and the anion (also the order which they are written)

NaI = sodium iodide

NH_4Cl = ammonium chloride

FeCl_2 = iron(II) chloride

FeCl = iron(I) chloride

Note: the Roman numerals are used as appropriate

3. Binary Molecular Compounds

The combination of two non-metals usually forms a binary compound. They usually contain two-worded names

The first word corresponds to the first element in the formula, with a Greek prefix to show the number of atoms of that element

2 = di 3 = tri 4 = tetra 5 = penta 6 = hexa
7 = hepta 8 = octa 9 = nona 10 = deca

The second word uses the stem name of the element, the Greek prefix, and the suffix **ide**


Examples: N_2O_5 = dinitrogen pentaoxide
 NO_2 = nitrogen dioxide
 N_2O = dinitrogen oxide

Many binary compounds have accepted common names:

H_2O = water NH_3 = ammonia
 H_2O_2 = hydrogen peroxide C_2H_2 = acetylene

4. Acids

Some binary compounds ionize in water to form H^+ ions, and are called **acids**.
For example, when hydrogen chloride is dissolved in water it forms H^+ and Cl^- ions.



The water solution of HCl is called hydrochloric acid

Acids that also contain oxygen are oxoacids.

Two common ones are nitric acid (HNO_3) and sulfuric acid (H_2SO_4)

The names of oxoacids originate from the names of the corresponding oxoanions.
The suffix **ate** is replaced with **ic**, and **ite** is replaced with **ous**

Examples:

ClO_4^- = perchlorate ion

$HClO_4$ = perchloric acid

ClO_3^- = chlorate ion

$HClO_3$ = chloric acid

ClO_2^- = chlorite ion

$HClO_2$ = chlorous acid


ClO^- = hypochlorite

$HClO$ = hypochlorous acid

Note: polyanions do not "break apart".

In the example below, sulfate stays as sulfate





The mole (M&H Ch. 3) (relevant problems found in tutorial manual section Stoichiometry Part 1, p. 17...)

The unit for chemical mass is the **mole**

≡ "the mass (in g) of an element or compound equal to its average atomic mass (in amu)"

▪ usually denoted by symbol "n"

For example 1 atom of C weighs 12.01 amu while 1 mole of C weighs 12.01 g

That methane, CH_4 : one molecule weighs $[12.01 + 4(1.01)] = 16.05$ amu while one mole weighs 16.05 g

One mole of a substance always contains the same number of particles
 $= 6.022 \times 10^{23}$
 ≡ Avogadro's number, N_{AV}

Compare 1 dozen = 12 units; 1 mole = 6.022×10^{23} units

For example: 1 mole of $\text{O}_2 = 6.022 \times 10^{23}$ O_2 molecules = 32.0 g

1 mole $\text{CH}_3\text{CH}_2\text{OH}$ contains 6.022×10^{23} ethanol molecules = 46.0 g
 and contains $9 \times N_{\text{AV}} = 5.42 \times 10^{24}$ atoms total
 of which $6 \times N_{\text{AV}} = 3.61 \times 10^{24}$ are hydrogen atoms, $2 \times N_{\text{AV}} = 1.20 \times 10^{24}$ are C atoms,
 6.022×10^{23} are O atoms

The **molar mass** of a molecule is the **sum** of the individual atomic masses.

For example the molar mass of $\text{CH}_3\text{CH}_2\text{OH} = 12 + 3 + 12 + 2 + 16 + 1 = 46.0 \text{ g mol}^{-1}$

Molar mass also termed the **molecular weight** or for ionic compounds such as NaCl , the **formula weight**

Worked Example:

If coal is 3.00 % by mass sulphur, S, how many moles of S are contained in 1 tonne (1000 kg) of coal?

Mass of S in 1 tonne of coal = $(3.00/100.0) \times 1000 \text{ kg} = 30.0 \text{ kg}$

$30 \text{ kg} = 30.0 \times 10^3 \text{ g}$. ∴ The number of moles, $n = (30.0 \times 10^3 \text{ g})/32 \text{ g mol}^{-1}$

$$n = 940 \text{ mol tonne}^{-1}$$

Molecular Formulas

Given a molecular formula, one can calculate the % composition of the elements present:

Worked Example:

for ethanol, $\text{CH}_3\text{CH}_2\text{OH} = \text{C}_2\text{H}_6\text{O}$ the molar mass = 46 g mol^{-1}

Mass of C in compound = $12 \times 2 = 24$

$$\therefore \% \text{C} = (24.0/46.0) \times 100\% = 52.2\%$$

Similarly, $\% \text{H} = ([6 \times 1.01]/46.0) \times 100\% = 13.2\%$

and

$$\% \text{O} = (16.0/46.0) \times 100\% = 34.8\%$$